

## Proposed Fish Stocking 2017

### Brown Trout



Stocking Location	Amount	Length	Weight
Saugeen River in Hanover (below dam)	2500	25-30 cm (10-12")	500 g (1 lb)
Allan Park Pond	2500	25-30 cm (10-12")	500 g (1 lb)

### Brook Trout



Stocking Location	Amount	Length	Weight
Bells Lake	3000	30 cm (12")	500 g (1 lb)
Feversham Pond	1500	30 cm (12")	500 g (1 lb)
Durham Town Pond	1500	30 cm (12")	500 g (1 lb)
Flesherton Town Pond	2000	30 cm (12")	500 g (1 lb)
Holstein Town Pond	2000	30 cm (12")	500 g (1 lb)

#### Stocking Strategy: Put and Take

This initiative focuses on MNRF stocking trout in local lakes and ponds with the expectation that anglers catch and keep the fish.

Trout will be stocked only days before the season opener in late April. Since these fish have spent their entire life cycle at the *Chatsworth Fish Culture Station*, they may not immediately recognize wild food sources. Use light 4-6 lbs. Line and keep hooks and bait small to mimic fish pellets that these fish are accustomed to. Small pieces of worm, power baits and small spinners (various colours) are the best bets. Also, try your artificial flies as they can sometimes out-fish anything else in the tackle box for these trout!

### Walleye



Stocking Location	Amount	Length	Weight
Chesley Lake	5000	15-23 cm (6-9")	

#### Stocking Strategy: Put—Grow—Take

This initiative focuses on MNRF stocking walleye in the fall, allowing them to grow to a suitable size, and anglers later catching and keeping these fish.

These fish are stocked in the late fall and will supplement the existing populations. Walleye are light sensitive and fishing is best early or late in the day. Look for walleye to move on to shoals or near structure to feed. A hook and weight, or jig-head tipped with a live minnow or work can be productive. Trolling slow with a worm harness and body bait lures can also be a good way to locate walleye.